

confidence interval (CI): 16.3–25.2%). Male sex (prevalence ratio (PR) 1.61; 95%CI 1.17–2.21), age \geq 80 years (PR 4.15; 95%CI 2.65–6.50), underweight (PR 1.90; 95%CI 1.23–2.93), diabetes mellitus (PR 1.80; 95%CI 1.20–2.73), and hospitalization in the past year (PR 2.00; 95%CI 1.36–2.91) were significantly associated with probable sarcopenia.

Keywords:

Brazil, EWGSOP2, older adults, rural population, sarcopenia.

FULL ARTICLE:

Introduction

The accelerated process of population aging has gained evidence in Brazil and other low- and middle-income countries¹. The increasing number of older adults in these countries has demanded greater care to prevent diseases and maintain the health of aging populations^{1,2}. In rural Brazil, population aging is characterized by higher proportion of men than women^{3,4}, low income³, precarious living^{5,6} and limited access to health care⁷.

Sarcopenia is related to the aging process and leads to a decline in the musculoskeletal system, leading to adverse health outcomes and higher health system costs⁸. One of the main research groups in this field, the European Working Group on Sarcopenia in Older People (EWGSOP), recommended the use of three different combinations of muscle assessment to diagnose sarcopenia (muscle mass, strength, and performance), with low muscle mass being the main criterion for diagnosing the disease⁹. An update made by EWGSOP2⁸, in 2019, recognized low muscle strength as the main criterion for identifying probable sarcopenia, and, when considered together with low muscle mass, diagnosis of the disease is confirmed. In addition, the group advised that, in order to assess the severity of sarcopenia, muscle performance should be evaluated, considering sarcopenia to be severe when muscle performance is low⁸.

The group⁸ also recommends using the Find-Access-Confirm-Severity (FACS) algorithm to identify and confirm sarcopenia cases. Screening begins with the Strength, Assistance with walking, Rising from a Chair, Climbing stairs, and Falls (SARC-F) questionnaire, followed by the analysis of the other criteria⁸. However, the validity of the SARC-F questionnaire has been questioned in the literature¹⁰. EWGSOP2 recognized its low sensitivity and moderate–high specificity, detecting primarily severe cases⁸. Recently, a meta-analysis demonstrated the inappropriateness of this test for sarcopenia screening¹⁰.

Recent updates and inconsistencies in the identification of sarcopenia have impacted the prevalence of sarcopenia in Brazil and worldwide. According to the EWGSOP2 criteria, sarcopenia prevalence ranges from 0.8% to 28%^{11–14}. In Brazil, prevalence varies from 10.1% to 33.7%^{15,16} and, when other assessment criteria are used, prevalence varies from 1.9% to 33.3%^{17–19}.

Despite these differences, the impact of sarcopenia is considerable^{8,9,15,20}. Falls, fractures, functional disability, hospitalization, and mortality are some of the factors associated with sarcopenia^{8,9,20}.

Considering the accelerated ageing process and the limited knowledge about the burden of sarcopenia and its consequences among older adults living in rural areas, this study aimed to estimate the prevalence of probable sarcopenia and its associated

Conclusion: One in five rural older adults have probable sarcopenia. Older adults living in rural areas need frequent measurements of muscle strength, and greater investments should be made to minimize the physical disability caused by sarcopenia among this population.

factors using EWGSOP2 criteria.

Methods

This cross-sectional study using data from a prospective cohort, the EpiRural Cohort Study, aimed to describe and monitor patterns of morbidity and mortality and the use of health services by older people (aged \geq 60 years) living in the rural area of the municipality of Rio Grande, in Rio Grande do Sul State, Brazil. Rio Grande is a municipality located in southern Brazil and, in 2017, its population was estimated as being 209 378 inhabitants, of whom approximately 5% lived in rural areas^{2,3}. The rural area of Rio Grande is delimited by the Laguna dos Patos, the Atlantic Ocean and Laguna Mirim water bodies, where traditional communities live mainly from fishing and family farming, including livestock, rice and soybean. The baseline data were collected in 2017, when 1029 older adults were identified through mapping and a random selection of permanently inhabited households in the rural area of the municipality. The first follow-up was conducted between 2018 and 2019 with the participation of 862 older adults (available from authors upon request). Further information about the EpiRural Cohort Study can be found elsewhere²¹.

The EpiRural's second follow-up was conducted between November 2020 and January 2022 (with unpredicted interruptions due to the COVID-19 pandemic). Two structured questionnaires were applied through face-to-face home interviews conducted by trained interviewers, accompanied by a field supervisor. Information was collected about sociodemographic, family, behavioral, and health characteristics of the older adults. Data were collected and managed using REDCap (Research Electronic Data Capture) electronic data capture tools hosted at the Federal University of Rio Grande²².

Probable sarcopenia was assessed using a handgrip strength test (HGS) performed with a Saehan hydraulic dynamometer, following the current standards^{8,23,24}. The HGS test was performed with the older adults seated in an armless chair without a back support, legs together, and feet on the floor, with the elbows in a position close to the body at 90° of flexion²⁴. In this position, the dynamometer was handed over to the participants and, after adjustment to the correct hand size, the older adults were asked to squeeze the equipment for 5 s with maximum force. The other hand remained in a neutral and relaxed position, resting on the opposite leg to which the HGS test was being applied. Three measurements were taken for each hand, starting with the right and then the left hand, with an interval of approximately 30 s of rest between each measurement²⁵. For analytical purposes, the highest measurement of the six was considered. The dependent variable, probable sarcopenia, was operationalized as the presence of low HGS, according to EWGSOP2⁸. The cut-off point for low muscle strength is HGS <27 kg for men and <16 kg for women^{8,26}.

The following independent variables were examined: sex (male/female), age (60–69 years; 70–79 years; ≥ 80 years), body mass index (BMI) (underweight: ≤ 22 kg/m²; healthy weight: 22–27 kg/m²; overweight: ≥ 27 kg/m²)²⁷, smoking status (never smoked; current smoker; former smoker), presence of self-reported diseases (hypertension (no/yes); diabetes mellitus (DM) (no/yes); chronic obstructive pulmonary disease (COPD) [no/yes) that was considered as the sum of the following self-reported diseases: asthma, bronchitis, emphysema; depression (measured with the Patient Health Questionnaire-9 (PHQ-9) (no/yes)), presence of spinal pain (no/yes) (considered as the sum of pain in the cervical, thoracic and lumbar regions of the spine), presence of knee pain (no/yes), and hospitalization in the previous 12 months (no/yes).

Data collection during the COVID-19 pandemic was performed following all social and sanitary norms.

Initially, a descriptive analysis of the data was conducted to identify the main characteristics of the older adults. The Shapiro–Wilk test was performed to assess the distribution of continuous data, and the symmetry of the distribution. Continuous variables were tested using the Mann–Whitney *U*-test. Prevalence of the outcome according to the independent variables was examined using the χ^2 test. Factors associated with probable sarcopenia were examined using crude and adjusted Poisson regression analysis. The adjusted analysis followed a hierarchical model controlling for confounding factors: first level (sex and age), second level (BMI and smoking status), third level (self-reported diagnosis of hypertension, DM, depression, COPD, presence of back pain, and presence of knee pain), and fourth level (hospitalization in the previous year). Variables with $p < 0.20$ were kept in the model and a 5% significance level was adopted.

Ethics approval

This research was approved by the Research Ethics Committee of

the Health Area of the Federal University of Rio Grande through Opinions No. 51/2017 and No. 154/2018. All ethical aspects were observed in accordance with National Health Council Resolution No. 466/12. Participation was voluntary, and all participants provided free, prior and informed consent.

Results

A total of 651 older adults were followed up in the second wave of the EpiRural Cohort Study in Rio Grande. We found that 51.9% were male, 45.8% were aged 70–79 years, and the majority (50.7%) were overweight (BMI ≥ 27 kg/m²). Among the diseases investigated, 59.7% of the participants self-reported hypertension, 15.0% DM, 17.4% COPD, and 7.1% depression (Table 1).

Prevalence of probable sarcopenia was 19.2% (95% confidence interval (CI) 16.3–22.5%). Occurrence of probable sarcopenia was higher among males (23.8% v 14.3%) and increased with age, reaching 44.3% in adults aged ≥ 80 years. Prevalence of probable sarcopenia was higher among older adults with diabetes (28.6%), depression (30.9%) and those who had been hospitalized in the previous year (41.4%) (Table 2).

Mean HGS was 27.8 kg (standard deviation (SD) ± 9.33), and males had a higher average (33.1 ± 8.58 kg) than females (22.1 ± 6.25 kg) (Table 3).

In the adjusted analysis, there was a linear gradient between age and probable sarcopenia, with those aged ≥ 80 years having a 4.15 times greater risk higher compared to those aged 60–70 years (prevalance ratio (PR) 4.15; 95%CI 2.65–6.50). In addition, there was association with the male sex (PR 1.61; 95%CI 1.17–2.21), underweight (BMI ≤ 22 kg/m²) (PR 1.90; 95%CI 1.23–2.93) in comparison to healthy weight individuals, diabetes mellitus (PR 1.80; 95%CI 1.20–2.73), and hospitalization in the previous year (PR 2.00; 95%CI 1.36–2.91) (Table 4).

Table 1: Characteristics of a sample of older adults living in the rural area of Rio Grande do Sul State, Brazil, 2020–2022 (n=651)

Variable	n (%)
Sex (n=651)	
Male	338 (51.9)
Female	313 (48.1)
Age (in complete years) (n=651)	
60–69	208 (31.9)
70–79	298 (45.8)
≥80	145 (22.3)
Body mass index, kg/ m ² (n=562)	
≤22	69 (12.3)
22–27	208 (37.0)
>27	285 (50.7)
Tobacco smoking (n=621)	
Never smoked	332 (53.5)
Current smoker	69 (11.1)
Former smoker	220 (35.4)
Hypertension (n=620)	
No	250 (40.3)
Yes	370 (59.7)
Diabetes mellitus (n=620)	
No	527 (85.0)
Yes	93 (15.0)
COPD (n=649)	
No	536 (82.6)
Yes	113 (17.4)
Depression (PHQ-9) (n=616)	
No	572 (92.9)
Yes	44 (7.1)
Spinal pain [†] (n=650)	
No	363 (55.8)
Yes	287 (44.2)
Knee pain (n=650)	
No	468 (72.0)
Yes	182 (28.0)
Hospitalization in the past year (n=651)	
No	617 (94.8)
Yes	34 (5.2)

[†] Sum of pain located in the cervical, thoracic, and lumbar regions of the spine. COPD, chronic obstructive pulmonary disease (emphysema/asthma/bronchitis). PHQ-9, Patient Health Questionnaire-9.

Table 2: Prevalence of probable sarcopenia, according to independent variables, in the rural area of Rio Grande do Sul State, Brazil, 2020–2022 (n=624)

Variable	Probable sarcopenia (EWGSOP2)	
	% (95%CI)	p-value (χ ² test)
Probable sarcopenia	19.2 (16.3–22.5)	–
Sex		0.003
Male	23.8 (19.4–28.7)	
Female	14.3 (10.8–18.8)	
Age (in complete years)		<0.001
60–69	10.4 (6.8–15.5)	
70–79	14.1 (10.5–18.6)	
≥80	44.3 (35.9–53.0)	
Body mass index, kg/m ²		<0.001
22	38.2 (27.3–50.5)	
22–27	16.0 (11.6–21.7)	
>27	15.5 (11.7–20.2)	
Tobacco smoking		0.817
Never smoked	19.1 (15.1–23.8)	
Current smoker	18.2 (10.5–29.7)	
Former smoker	21.0 (16.1–27.2)	
Hypertension		0.618
No	18.7 (14.20–24.3)	
Yes	20.4 (16.5–24.9)	
Diabetes mellitus		0.021
No	18.1 (15.0–21.7)	
Yes	28.6 (20.1–38.9)	
Depression (PHQ-9)		0.026
No	17.2 (14.3–20.5)	
Yes	30.9 (18.4–47.0)	
COPD		0.414
No	19.6 (16.3–23.2)	
Yes	16.2 (10.4–24.4)	
Spinal pain [†]		0.130
No	21.4 (17.4–26.2)	
Yes	16.6 (12.7–21.4)	
Knee pain		0.054
No	17.3 (14.0–21.2)	
Yes	24.2 (18.3–30.9)	
Hospitalization in the past year		0.002
No	18.1 (15.3–21.5)	
Yes	41.4 (24.3–60.7)	

[†] Sum of pain located in the cervical, thoracic, and lumbar regions of the spine.
 CI, confidence interval. COPD, chronic obstructive pulmonary disease (emphysema/asthma/bronchitis).
 EWGSOP, European Working Group on Sarcopenia in Older People. PHQ-9, Patient Health Questionnaire-9.

Table 3: Mean, standard deviation and median measurements of handgrip strength, calf circumference and gait speed of older adults living in the rural area of Rio Grande do Sul State, Brazil, 2020–2022

Variable	All		Male		Female		p-value (Mann-Whitney U-test)
	Mean±SD	Median	Mean±SD	Median	Mean±SD	Median	
Handgrip strength test, kg	27.8±9.33	26.8	33.1±8.58	33.6	22.1±6.25	21.3	<0.001

SD, standard deviation.

Table 4: Crude and adjusted analysis of associations between prevalence of probable sarcopenia (EWGSOP2) and independent variables, rural area of Rio Grande do Sul State, Brazil, 2020–2022

Variable	Probable sarcopenia (EWGSOP2)	
	Crude analysis PR (95%CI)	Adjusted analysis PR (95%CI)
Sex	<i>p</i> =0.003*	<i>p</i> =0.005*
Male	1.65 (1.18–2.32)	1.61 (1.17–2.21)
Female	1.00	1.00
Age (in complete years)	<i>p</i> <0.001**	<i>p</i> <0.001**
60–69	1.00	1.00
70–79	1.35 (0.82–2.22)	1.31 (0.80–2.15)
≥80	4.25 (2.71–6.67)	4.15 (2.65–6.50)
Body mass index, kg/m ²	<i>p</i> <0.001*	<i>p</i> =0.011*
≤22	2.38 (1.54–3.69)	1.90 (1.23–2.93)
22–27	1.00	1.00
>27	0.97 (0.63–1.46)	1.32 (0.87–2.00)
Tobacco smoking	<i>p</i> =0.816*	<i>p</i> =0.623*
Never smoked	1.00	1.00
Current smoker	0.95 (0.54–1.66)	0.78 (0.40–1.50)
Former smoker	1.10 (0.78–1.55)	0.85 (0.58–1.25)
Hypertension	<i>p</i> =0.618*	<i>p</i> =0.985*
No	1.00	1.00
Yes	1.09 (0.78–1.53)	1.00 (0.70–1.44)
Diabetes mellitus	<i>p</i> =0.017*	<i>p</i> =0.006*
No	1.00	1.00
Yes	1.57 (1.08–2.30)	1.80 (1.20–2.73)
Depression (PHQ-9)	<i>p</i> =0.018*	<i>p</i> =0.066*
No	1.00	1.00
Yes	1.80 (1.10–2.92)	1.57 (0.97–2.54)
COPD	<i>p</i> =0.421*	<i>p</i> =0.756*
No	1.00	1.00
Yes	0.82 (0.54–1.31)	0.92 (0.58–1.50)
Spinal pain [†]	<i>p</i> =0.133*	<i>p</i> =0.161*
No	1.00	1.00
Yes	0.78 (0.55–1.08)	0.78 (0.55–1.10)
Knee pain	<i>p</i> =0.051*	<i>p</i> =0.148*
No	1.00	1.00
Yes	1.39 (0.99–1.93)	1.28 (0.91–1.80)
Hospitalization in the past year	<i>p</i> <0.001*	<i>p</i> =0.001*
No	1.00	1.00
Yes	2.28 (1.43–3.63)	2.00 (1.36–2.91)

* Heterogeneity test

** Wald trend test

[†] Sum of pain located in the cervical, thoracic, and lumbar regions of the spine.

CI, confidence interval. COPD, chronic obstructive pulmonary disease (emphysema/asthma/bronchitis).

EWGSOP, European Working Group on Sarcopenia in Older People. PHQ-9, Patient Health Questionnaire-9. PR, prevalence ratio.

Discussion

This study found that one in five older adults living in rural areas in the far south of Brazil has probable sarcopenia. Male sex, age ≥80 years, underweight, having DM and having been hospitalized in the previous year were associated with the outcome.

In Brazil, few population-based studies have assessed probable sarcopenia and only one reported similar prevalence to our findings¹⁵. Similar results were observed in Asia in a study of older adults living in urban and rural areas¹⁴. In Europe, prevalence of probable sarcopenia is higher than that reported in Brazil^{11,15,28,29}. There are few studies analyzing the differences in the occurrence of probable sarcopenia between areas of residence (urban v rural). However, rural prevalence of sarcopenia in older adults has great variability^{30–32} and these divergences may be related to recent updates of sarcopenia criteria.

One unexpected finding of our study was that males have a higher risk of probable sarcopenia than females, contradicting previous studies suggesting that females in Latin America are at a greater risk of sarcopenia³³, including a Peruvian study carried out only with older adults living in rural areas³². In Portugal, protection for males was identified using the EWGSOP2 criteria for diagnosis of sarcopenia (odds ratio (OR) 0.52 (0.29–0.97))²⁹.

Another Brazilian study found similar results but using different diagnosis criteria¹⁹. In southern Iran, males are also at higher risk of sarcopenia (OR 3.13 (91.23–7.98))³⁴. This may possibly be due to the high cut-off point indicated by EWGSOP2 to consider low muscle strength in men, which is much higher than that applied in

women, leading to an overestimation of probable sarcopenia in men. A recent Brazilian study suggested a lower HGS cutoff for men³⁵, which may be more reliable in relation to the Brazilian population. Therefore, new studies with other HGS cut-off points for males are needed to confirm or refute this association. Another possible explanation is the fact that male are more likely to develop work-related injuries in rural and fishing activities.

Association between aging and probable sarcopenia has also been found in studies that used the updated EWGSOP2 criteria^{12,29} and studies using several other criteria^{19,30,32,33}. This can be explained by the natural process of body aging that involves increased protein degradation, changes in body composition, increased fat infiltration into muscle tissue, increased inflammatory cytokines, decreased body homeostasis, and decreased circulating hormones, which contributes to the risk of sarcopenia^{36,37}. Notably, older people aged 70–79 years were not at risk of probable sarcopenia. This finding can be explained by the fact that some older adults remain active and perform activities that require greater physical effort on their rural properties, with less sedentary behavior when compared to older adults living in urban areas³⁸. Another possible explanation can be the peculiarity verified in the most recent Brazilian agricultural census (2017)³, in which 23% of rural producers were ≥65 years old³, suggesting that a large proportion of Brazilian older adult residents in these locations participate in the productive scenario in the Brazilian countryside.

Several studies conducted in Brazil and worldwide have reported association between underweight and sarcopenia^{14,19,30,34,39}. The BMI measurement is easily performed in both clinical practice and population-based studies, and is considered a reliable indicator of

nutritional status in older adults^{27,40}. Although there are differences in the cut-off points for BMI in older adults^{40,41}, the literature is consistent with the assumption that underweight in the older adults should be monitored frequently, as it increases the risk of malnutrition, and consequently the impairment of health, increasing the risk of sarcopenia, frailty, vulnerability, and mortality⁴¹.

The association of diabetes mellitus with probable sarcopenia observed in our study (PR 1.89 (1.30–2.75)) was also found in studies performed in Asia⁴² and Europe⁴³. A recent meta-analysis of observational studies found that one disease leads to another, especially in individuals who already have DM complications⁴⁴. Some of the pathophysiological mechanisms of DM also occur in sarcopenia, including increased insulin resistance, chronic inflammation, and mitochondrial changes⁴⁵. In addition, Park et al (2006)⁴⁶ identified that older adults with DM have lower muscle strength. However, our research did not analyze the type of DM (type I or II), time of disease onset, or use of specific medications, which are suggestions for future research.

Association between probable sarcopenia and hospitalization in the previous year is well established in the literature^{8,20,47}. Hospitalization leads to decreased muscle strength and mobility, mainly because of the negative effects of a long stay in hospital beds, which can increase the risk of sarcopenia among the older adults, especially those affected by comorbidities^{43,47,48}. Thus, hospitalization is an important risk factor, suggesting the need for intervention during and after this period, increasing body mobility and encouraging performance of guided exercises.

However, this study had some limitations. Its cross-sectional design presents a reverse causality bias, which makes it impossible to establish a cause-and-effect mechanism between the associations, especially in relation to BMI. Additionally, information on level of physical activity, cognitive impairment and quality of life in the older adult participants was not collected, making this analysis unfeasible. Lastly, some health information may be biased since it was self-reported.

Among its strengths, our study followed the EWGSOP2 updates to diagnose probable sarcopenia, being, to the best of our knowledge, the first Brazilian population-based study to measure the occurrence of sarcopenia by applying these criteria to older adults living in a rural area. We believe that EWGSOP2 enhances our comparability with the literature. In addition, the easy use of a dynamometer, and its low cost, make it a feasible resource for clinical practice of primary health care, facilitating access to screening probable sarcopenia, and initiating necessary interventions as early as possible^{8,49,50}. We suggest further research on sarcopenia in rural populations.

Conclusion

Probable sarcopenia can easily be identified in primary health care and reversed through specific health care. Older adults living in rural areas require especial attention, particularly males, individuals aged ≥ 80 years, underweight, diabetics and those hospitalized in the previous year.

In addition, we highlight the importance of having more health professionals working directly in musculoskeletal health care, as well as investment in facilities for health promotion and engagement in resistance exercise programs, allied with changes in lifestyle that can support reversing the condition of probable sarcopenia and minimize its impact on older adults' health.

Finally, we believe that our findings can contribute to directing public health policies for older adults by investing in the prevention of probable sarcopenia.

Funding

The authors acknowledge the Research Support Foundation of Rio Grande do Sul (FAPERGS).

Conflicts of interest

The authors declare that there is no conflict of interest.

REFERENCES:

- 1 Pan American Health Organization (PAHO)/World Health Organization (WHO). Decade of Healthy Aging (2021–2030) in the Americas. [In Portuguese]. *More60 Studies on Aging* 2021; **32**: 80. Available: [web link](#) (Accessed 13 April 2022).
- 2 Brazilian Institute of Geography and Statistics – IBGE. *Projections of the population of Brazil and the Federation Units*. [In Portuguese]. 2019. Available: [web link](#) (Accessed 12 May 2022).
- 3 Brazilian Institute of Geography and Statistics – IBGE. *Agricultural Census - definitive results*. [In Portuguese]. 2017. Available: [web link](#) (Accessed 6 May 2022).
- 4 Froehlich JM, Rauber CC, Carpes RH, Toebe M. Selective exodus, masculinization and rural aging in the central region of the RS. *Sociologia Rural. Ciência Rural* 2011; **41(9)**: 1674–1680. DOI link
- 5 Soares RAS, Moraes EM, Viana RPT, Pessoa VM, Carneiro FF. Social environmental determinants and health: rural Brazil versus Brazil urban. *Tempus Actas de Saúde Coletiva* 2015; **9**: 221–235. DOI link
- 6 Castilho EA, Gonçalves H. Health problems and the rural area. *Revista de Saúde Pública* 2018; **52**: 1s. DOI link
- 7 Magalhães DL, Matos RS, Souza AO, Neves RF, Costa MMB, Rodrigues AA, et al. Access to health and quality of life in the rural area. *Research, Society and Development* 2022; **11**: e50411326906. DOI link
- 8 Cruz-Jentoft AJ, Bahat G, Bauer J, Boirie Y, Bruyère O, Cederholm T, et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age and Ageing* 2019; **48(1)**: 16–31. DOI link, PMID:30312372
- 9 Cruz-Jentoft AJ, Baeyens JP, Bauer JM, Boirie Y, Cederholm T, Landi F, et al. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age and Ageing* 2010; **39(4)**: 412–423. DOI link, PMID:20392703
- 10 Voelker SN, Michalopoulos N, Maier AB, Reijnierse EM. Reliability and concurrent validity of the SARC-F and its modified versions: a systematic review and meta-analysis. *Journal of the American Medical Directors Association* 2021; **22(9)**: 1864–1876.e16.
- 11 Lampignano L, Bortone I, Castellana F, Donghia R, Guerra V, Zupo R, et al. Impact of different operational definitions of

- sarcopenia on prevalence in a population-based sample: The Salus in Apulia Study. *International Journal of Environmental Research and Public Health* 2021; **18(24)**: 12979. DOI link, PMID:34948590
- 12** Wearing J, Konings P, Bie RA, Stokes M, Bruin ED. Prevalence of probable sarcopenia in community-dwelling older Swiss people - a cross-sectional study. *BMC Geriatrics* 2020; **20(1)**: 307. DOI link, PMID:32847545
- 13** Wu X, Li X, Xu M, Zhang Z, He L, Li Y. Sarcopenia prevalence and associated factors among older Chinese population: Findings from the China Health and Retirement Longitudinal Study. *PLoS One* 2021; **16(3)**: e0247617. DOI link, PMID:33661964
- 14** Kim M, Won CW. Prevalence of sarcopenia in community-dwelling older adults using the definition of the European Working Group on Sarcopenia in Older People 2: findings from the Korean Frailty and Aging Cohort Study. *Age and Ageing* 2019; **48(6)**: 910-916. DOI link, PMID:31329815
- 15** Bachettini NP, Bielemann RM, Barbosa-Silva TG, Menezes A, Tomasi E, Gonzalez MC. Sarcopenia as a mortality predictor in community-dwelling older adults: a comparison of the diagnostic criteria of the European Working Group on Sarcopenia in Older People. *European Journal of Clinical Nutrition* 2020; **74(4)**: 573-580. DOI link, PMID:31586126
- 16** Marincolo J, Aprahamian I, Corona LP, Neri AL, Yassuda MS, Borim F. Three definitions of probable sarcopenia and associations with falls and functional disability among community-dwelling older adults. *Osteoporosis and Sarcopenia* 2021; **7(2)**: 69-74. DOI link, PMID:34278002
- 17** Barbosa-Silva TG, Bielemann RM, Gonzalez MC, Menezes AM. Prevalence of sarcopenia among community-dwelling elderly of a medium-sized South American city: results of the COMO VAI? Study. *Journal of Cachexia, Sarcopenia and Muscle* 2016; **7(2)**: 136-143. DOI link, PMID:27493867
- 18** Szlejf C, Suemoto CK, Brunoni AR, Viana MC, Moreno AB, Matos SMA, et al. Depression is associated with sarcopenia due to low muscle strength: results from the ELSA-Brasil Study. *Journal of the American Medical Directors Association* 2019; **20(12)**: 1641-1646. DOI link, PMID:30409492
- 19** Pelegrini A, Mazo GZ, Pinto AA, Benedetti T, Silva DAS, Petroski EL. Sarcopenia: prevalence and associated factors among elderly from a Brazilian capital. *Fisioterapia em Movimento* 2018; **31**: e003102. DOI link
- 20** Beaudart C, Zaaria M, Pasleau F, Reginster JY, Bruyère O. Health outcomes of sarcopenia: a systematic review and meta-analysis. *PLoS One* 2017; **12(1)**: e0169548. DOI link, PMID:28095426
- 21** Meucci RD, Farias CP, Paludo CS, Pagliaro G, Soares MP, Lima SH, et al. Aging in a rural area in southern Brazil: designing a prospective cohort study. *Rural and Remote Health* 2022; **22(1)**: 6591. DOI link, PMID:35192773
- 22** Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap) - a metadata-driven methodology and workflow process for providing translational research informatics support. *Journal of Biomedical Informatics* 2009; **42(2)**: 377-381. DOI link, PMID:18929686
- 23** Habicht JP. Standardization of quantitative epidemiological methods in the field. *Boletín de la Oficina Sanitaria Panamericana* 1974; **76(5)**: 375-384. PMID:4277063
- 24** Fess EE. *Grip strength*. 2nd edn. Chicago: American Society of Hand Therapists, 1992.
- 25** Roberts HC, Denison HJ, Martin HJ, Patel HP, Syddall H, Cooper C, et al. A review of the measurement of grip strength in clinical and epidemiological studies: towards a standardised approach. *Age and Ageing* 2011; **40(4)**: 423-429. DOI link, PMID:21624928
- 26** Dodds RM, Syddall HE, Cooper R, Benzeval M, Deary IJ, Dennison EM, et al. Grip strength across the life course: normative data from twelve British studies. *PLoS One* 2014; **9(12)**: e113637. DOI link, PMID:25474696
- 27** Brazil Ministry of Health, Department of Health Care, Department of Primary Care. *Guidelines for the collection and analysis of anthropometric data in health services: Technical Standard of the Food and Nutrition Surveillance System - SISVAN*. [In Portuguese]. 2011. Available: [web link](#) (Accessed 13 October 2022).
- 28** Tsekoura M, Billis E, Matzaroglou C, Tsepis E, Gliatis J. Prevalence of probable sarcopenia in community-dwelling older Greek people. *Journal of Frailty, Sarcopenia and Falls* 2021; **6(4)**: 204-208. DOI link, PMID:34950810
- 29** Sousa-Santos AR, Afonso C, Borges N, Santos A, Padrão P, Moreira P, et al. Factors associated with sarcopenia and undernutrition in older adults. *Nutrition & Dietetics* 2019; **76(5)**: 604-612. DOI link, PMID:31083820
- 30** Gao L, Jiang J, Yang M, Hao Q, Luo L, Dong B. Prevalence of sarcopenia and associated factors in Chinese community-dwelling elderly: comparison between rural and urban areas. *Journal of the American Medical Directors Association* 2015; **16(11)**: 1003.e1-1003.e10036. DOI link, PMID:26385304
- 31** Wu X, Li X, Xu M, Zhang Z, He L, Li Y. Sarcopenia prevalence and associated factors among older Chinese population: Findings from the China Health and Retirement Longitudinal Study. *PLoS One* 2021; **16(3)**: e0247617. DOI link, PMID:33661964
- 32** Tramontano A, Veronese N, Sergi, G, Manzato E, Rodriguez-Hurtado D, Maggi S, et al. Prevalence of sarcopenia and associated factors in the healthy older adults of the Peruvian Andes. *Archives of Gerontology and Geriatrics* 2017; **68**: 49-54. DOI link, PMID:27649513
- 33** Samper-Ternent R, Reyes-Ortiz C, Ottenbacher KJ, Cano CA. Frailty and sarcopenia in Bogotá: results from the SABE Bogotá Study. *Ageing Clinical and Experimental Research* 2017; **29(2)**: 265-272. DOI link, PMID:27034288
- 34** Nasimi N, Dabbaghmanesh MH, Sohrabi Z. Nutritional status and body fat mass: Determinants of sarcopenia in community-dwelling older adults. *Experimental Gerontology* 2019; **122**: 67-73. DOI link, PMID:31022445
- 35** Fernandes S, Lima de Andrade LE, Dos Santos Aguiar Gonçalves RS, Aires da Câmara SM, Guerra RO, Cavalcanti Maciel AC. Cut-off points to screening for sarcopenia in community-dwelling older people residents in Brazil. *PeerJ* 2021; **9**: e12038. DOI link, PMID:34527442
- 36** da Costa JP, Vitorino R, Silva GM, Vogel C, Duarte AC, Rocha-Santos T. A synopsis on aging - Theories, mechanisms and future prospects. *Ageing Research Reviews* 2016; **29**: 90-112. DOI link, PMID:27353257
- 37** Shimokata H, Shimada H, Satake S, Endo N, Shibasaki K, Ogawa S, et al. Chapter 2. Epidemiology of sarcopenia. *Geriatrics & Gerontology International* 2018; **18**: 13-22. DOI link,

PMid:29745460

38 Leão OAA, Knuth AG, Meucci RD. Sedentary behavior in elderly residents from the rural area in Southern Brazil. *Brazilian Journal of Epidemiology* 2020; **23**: e200008. DOI link, PMid:32130397

39 Nunes JD, Zacarin JF, Pavarini SCI, Zazzetta MS, Orlandi AAS, Orlandi FS. Factors associated with sarcopenia in elderly community. [In Portuguese]. *Fisioterapia e Pesquisa* 2021; **28**(2): 159-165. DOI link

40 Francisco PMSB, Assumpção D, Borim FSA, Bacurau AGM, Malta DC. Underweight in older adults according to age and cutoff points of Body Mass Index. *Cadernos Saúde Coletiva* 2021; **29**(2): 246-250. DOI link

41 Winter JE, MacInnis RJ, Wattanapenpaiboon N, Nowson CA. BMI and all-cause mortality in older adults: a meta-analysis. *The American Journal of Clinical Nutrition* 2014; **99**(4): 875-890. DOI link, PMid:24452240

42 Su Y, Hirayama K, Han TF, Izutsu M, Yuki M. Sarcopenia prevalence and risk factors among Japanese community dwelling older adults living in a snow-covered city according to EWGSOP2. *Journal of Clinical Medicine* 2019; **8**(3): 291. DOI link, PMid:30823497

43 Martone AM, Bianchi L, Abete P, Bellelli G, Bo M, Cherubini A, et al. The incidence of sarcopenia among hospitalized older patients: results from the Glisten study. *Journal of Cachexia, Sarcopenia and Muscle* 2017; **8**(6): 907-914. DOI link, PMid:28913934

44 Veronese N, Pizzol D, Demurtas J, Soysal P, Smith L, Sieber C, et al. Association between sarcopenia and diabetes: a systematic

review and meta-analysis of observational studies. *European Geriatric Medicine* 2019; **10**(5): 685-696. DOI link, PMid:34652699

45 Umegaki H. Sarcopenia and frailty in older patients with diabetes mellitus. *Geriatrics & Gerontology International* 2016; **16**(3): 293-299. DOI link, PMid:26799937

46 Park SW, Goodpaster BH, Strotmeyer ES, de Rekeneire N, Harris TB, Schwartz AV, et al. Decreased muscle strength and quality in older adults with type 2 diabetes: the health, aging, and body composition study. *Diabetes* 2006; **55**(6): 1813-1818. DOI link, PMid:16731847

47 Sobestiansky S, Michaelsson K, Cederholm T. Sarcopenia prevalence and associations with mortality and hospitalisation by various sarcopenia definitions in 85-89 year old community-dwelling men: a report from the ULSAM study. *BMC Geriatrics* 2019; **19**(1): 318. DOI link, PMid:31747923

48 Loyd C, Markland AD, Zhang Y, Fowler M, Harper S, Wright NC, et al. Prevalence of hospital-associated disability in older adults: a meta-analysis. *Journal of the American Medical Directors Association* 2020; **21**(4): 455-461.e5. DOI link, PMid:31734122

49 Steffl M, Bohannon RW, Sontakova L, Tufano JJ, Shiells K, Holmerova I. Relationship between sarcopenia and physical activity in older people: a systematic review and meta-analysis. *Clinical Interventions in Aging* 2017; **12**: 835-845. DOI link, PMid:28553092

50 Wang Z, Xu X, Gao S, Wu C, Song Q, Shi Z, et al. Effects of internet-based nutrition and exercise interventions on the prevention and treatment of sarcopenia in the elderly. *Nutrients* 2022; **14**(12): 2458. DOI link, PMid:35745187